



Viburnum acerifolium
mapleleaf viburnum

SHADY INVADERS

Why Observe?

Your observations will help us quantify the extent to which invasive shrubs exhibit extended leaf phenology (having leaves for a longer period of time than natives in the same location). The Shady Invaders campaign compares the leaf phenology of native (mapleleaf viburnum) and invasive shrubs across the eastern U.S. in order to improve the timing/planning of land management activities, to predict how research on the impacts of extended leaf phenology (ELP) at one location applies across the range, and to examine how climate change may influence ELP.

Description

Mapleleaf viburnum twigs have tiny hairs. The flowers are white and monoecious.

Tips for Identification

These shrubs are commonly mistaken for maple tree saplings. However, the leaves of viburnum are fuzzy, unlike most maple tree leaves, which tend to be hairless and even glossy.

Be aware there is variation from individual to individual within a species, so your plant may not look exactly like the one pictured. If you are uncertain whether or not a phenophase is occurring, report a “?” for its status until it becomes clear what you are observing after subsequent visits.



Erynn Maynard



Erynn Maynard



Image Needed

email
photos@usanpn.org

Breaking leaf buds

One or more breaking leaf buds are visible on the plant. A leaf bud is considered "breaking" once a green leaf tip is visible at the end of the bud, but before the first leaf from the bud has unfolded to expose the leaf stalk (petiole) or leaf base.



Leaves

One or more live, unfolded leaves are visible on the plant. A leaf is considered "unfolded" once its entire length has emerged from a breaking bud, stem node or growing stem tip, so that the leaf stalk (petiole) or leaf base is visible at its point of attachment to the stem. Do not include fully dried or dead leaves.



Increasing leaf size

A majority of leaves on the plant have not yet reached their full size and are still growing larger. Do not include new leaves that continue to emerge at the ends of elongating stems throughout the growing season.




Colored leaves

One or more leaves show some of their typical late-season color, or yellow or brown due to drought or other stresses. Do not include small spots of color due to minor leaf damage, or dieback on branches that have broken. Do not include fully dried or dead leaves that remain on the plant.



Flowers or flower buds

One or more fresh open or unopened flowers or flower buds are visible on the plant. Include flower buds or inflorescences that are swelling or expanding, but do not include those that are tightly closed and not actively growing (dormant). Also do not include wilted or dried flowers.



Open flowers

One or more open, fresh flowers are visible on the plant. Flowers are considered "open" when the reproductive parts (male stamens or female pistils) are visible between or within unfolded or open flower parts (petals, floral tubes or sepals). Do not include wilted or dried flowers.

Image Needed

email
photos@usanpn.org

Fruits

One or more fruits are visible on the plant. For *Viburnum acerifolium*, the fruit is berry-like and changes from green to red to blue-black, purplish-black or black.



Ripe fruits

One or more ripe fruits are visible on the plant. For *Viburnum acerifolium*, a fruit is considered ripe when it has turned blue-black, purplish-black or black.

Phenophases not pictured: Falling leaves, Recent fruit or seed drop